







Saudi Arabia–United States Collaboration in Health Research: A Formula for Success

T. I. Hooper T. C. Smith, G. C. Gray M. S. Al Qahtani, Z.A. Memish D. H. Barrett, K. M. Schlangen



Naval Health Research Center

.Report No. 04-12

Approved for public release; distribution unlimited.

Naval Health Research Center 140 Sylvester Road San Diego, California 92106

AJC practice forum

Saudi Arabia-United States collaboration in health research: A formula for success

Tomoko I. Hooper, MD, MPH,^a Tyler C. Smith, MS,^b Gregory C. Gray, MD, MPH,^c Mohammed S. Al Qahtani, MBBS, DFE,^d Ziad A. Memish, MD,^d Drue H. Barrett, PhD,^e Karen M. Schlangen, MA,^b David F. Cruess, PhD,^a Margaret A. K. Ryan, MD, MPH,^b and Gary D. Gackstetter, DVM, MPH, PhD^a Bethesda, Maryland

The aim of this article is to share our experiences from an international collaborative effort to study health outcomes among Saudi Arabian National Guard (SANG) soldiers following the 1991 Gulf War. By paying particular attention to distinct social and religious customs, geopolitical differences, and unique aspects of the health care system, we achieved a successful international collaboration in health research. (Am J Infect Control 2005;33:192-6.)

International collaborations offer new perspectives and important opportunities for health researchers to share information, experiences, and methods that can provide the basis for population-based health studies of mutual interest. Collaborations that involve completely different cultural settings also present unique challenges. We assembled a team of 10 researchers: 2 from the Kingdom of Saudi Arabia and 8 from the United States to collaborate on a study of health outcomes

From the Department of Preventive Medicine and Biometrics at the Uniformed Services University, Bethesda, Maryland^a; Department of Defense Center for Deployment Health Research, Naval Health Research Center, San Diego, California^b; Department of Epidemiology, College of Public Health, University of Iowa, Iowa City, Iowa^c; Infection Prevention and Control Program, King Abdul Aziz Medical City, King Fahad National Guard Hospital, National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia^d; and Office of the Director, National Center for Environmental Health, and Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention, Atlanta, Georgia.^e

Reprint requests: Tomoko Hooper, Uniformed Services University of the Health Sciences, Department of Preventive Medicine and Biometrics-Room A1040G, 4301 Jones Bridge Road, Bethesda, MD 20814-4799. E-mail: thooper@usuhs.mil.

The views expressed in this article are those of the authors and do not reflect the official position of the Uniformed Services University of the Health Sciences, the Department of the Air Force, the Department of the Navy, the Department of Defense, the Centers for Disease Control and Prevention, or the US Government nor the views of the Saudi Arabian National Guard, Health Affairs, Kingdom of Saudi Arabia.

This report is public domain and can be copied freely.

0196-6553/\$30.00

Copyright © 2005 by the Association for Professionals in Infection Control and Epidemiology, Inc.

doi:10.1016/j.ajic.2005.01.006

among Saudi Arabian National Guard (SANG) soldiers following the 1991 Gulf War. In addition to distinct geopolitical differences, these 2 nations have vastly differing social and religious customs, languages, and health care systems. In this article, we describe our experiences with a successful international research collaboration. Our study findings are reported elsewhere. In the successful international research collaboration.

Our research collaboration was initiated in response to talks between the Chief Executive Officer of the Saudi Arabian National Guard Health Affairs at King Abdul Aziz Medical City (KAMC), formerly known as King Fahad National Guard Hospital, based in Riyadh, Kingdom of Saudi Arabia; and the Assistant Secretary of Defense for Gulf War Illnesses, US Department of Defense (DOD). The purpose of our study was to complement the multinational, more than decadelong research endeavor that has attempted to characterize the morbidity reported among some veterans of the 1991 Gulf War. 2-38 To establish the feasibility of a Saudi-US research collaboration and to ensure support for the initial study proposal through the appropriate Saudi channels, a 4-member team of US researchers from the Uniformed Services University of the Health Sciences (USUHS), Bethesda, Maryland, and the Centers for Disease Control and Prevention (CDC), Atlanta, Georgia, as well as a representative from the Office of the Assistant Secretary of Defense for Gulf War Illnesses traveled to Saudi Arabia in late 1998. Data collection efforts subsequently began in earnest in 1999 when 8 investigators from the United States (USUHS, CDC, and the Naval Health Research Center, NHRC) joined the 2 principal Saudi researchers in Riyadh to inventory and evaluate multiple, potential sources of health data at

193

KAMC, as well as other health care facilities in the Riyadh area. The senior status and size of the research group lent credibility to the collaborative effort and gave impetus to the process of obtaining data-use authorization. Members of the research team had expertise in epidemiology, biostatistics, data programming and analysis, and database management, as well as familiarity with the health issues surrounding the 1991 Gulf War. 6,10-16,29-33 In addition to the small, existing US military presence in Riyadh, the CDC was a well-known entity at KAMC because of its previous sponsorship of in-country epidemiologic training. Saudi coinvestigators, based at KAMC, provided technical expertise and key liaison with professional staff and senior hospital and SANG military leadership. The diverse nature of the research team ensured positive interactions with clinicians, pharmacists, hospital administrators, and information systems managers at KAMC.

The context for the SANG health study revolved around the lingering concerns over multisymptom conditions reported by some US veterans deployed to the 1991 Gulf War, as well as similar self-reported symptoms among some of the other coalition forces (e.g., United Kingdom, Canada, and Australia). 3,5,6,12,15,23,35-38 The opportunity to explore the health impact of the Gulf War on a coalition force native to the Gulf region was timely and held the promise of adding a unique perspective to our growing knowledge base on the possible health effects of deployment. At the same time, the distinct features of this research setting posed a number of challenges. For purposes of discussion, we have grouped these challenges into 3 topic areas: (1) sociocultural (language, family structure, religious observances, role of women, and social customs); (2) geopolitical (political climate, governance, and geographic/environmental factors); and (3) health care systems (eligibility, information systems, utilization patterns, and patient confidentiality issues).

SOCIOCULTURAL CHALLENGES

As members of a Saudi-US research team, we were keenly aware of the differences between our 2 nations outside the domain of science. Some of these differences had the potential to affect our ability to access data, construct an analyzable data set, and interpret our findings. Saudi researchers on the team played an essential role in facilitating data collection by removing barriers that could have impeded progress on our study.

Language differences complicated all aspects of our study. For example, personnel data from SANG Headquarters were in the form of paper records written in Arabic. As a result, each data field required translation into English prior to being key entered into a database. Additionally, a large number of SANG members had exactly the same name because of the Saudi propensity to use certain given and surnames. Saudi clans or extended family members can potentially number in the millions, and perhaps hundreds in a village can have identical names. These and other language issues complicated the process of assembling appropriate study and comparison groups. On the other hand, both Saudi investigators on our research team spoke fluent English, thus enhancing our ability to discuss study objectives, data requirements, and analysis plans as we implemented the protocol.

Complete accounting of family members was also challenging because of notable differences in family structure. For example, it is not unusual for SANG enlisted members to have 2 or 3 wives and 10 or more children in their household.

As guests in the Kingdom, it was essential for US researchers to be considerate of religious customs, such as Islamic prayer times. The 3 US women on the research team had to adjust to the distinct role of women in a Muslim society and were appropriately respectful of local customs related to dress, religious observances, and restricted movement in public. For instance, women are only allowed in certain areas (family restaurants) for public dining or in designated seating areas in public places. In addition, Saudi women are prohibited from driving vehicles or being seen in public unless accompanied by a spouse or adult male relative. Our visit to SANG Headquarters to request personnel data was an extraordinary event because it included the entire research team, and women are generally not allowed within this compound. The US women dressed appropriately in longsleeved, full-length over-garments, called abeyas, and covered their heads. Senior SANG officials were gracious hosts, welcoming the entire group.

GEOPOLITICAL CHALLENGES

Travel to the Kingdom of Saudi Arabia by the team of US researchers was allowed only under official government orders and required high-level support from the US Office of the Assistant Secretary of Defense and Saudi Arabian National Guard Health Affairs. Entry into the Kingdom was facilitated by US military officers assigned to SANG. Travel plans were coordinated to avoid conflict with religious observances, such as Ramadan or Christmas, as well as to avoid extremes in the physical environment. As international travelers, we had to remain aware of current political sensitivities and global events that could affect our safety and our ability to travel freely.

Scheduling of teleconferences had to take into account at least 10 time zones, when the entire 194 Vol. 33 No. 3

research team was involved, to connect the Western US and points in-between to Riyadh, Saudi Arabia. In addition, meetings and teleconferences had to allow for differences in how the workweek is defined in Saudi Arabia and the United States. For Saudis, Thursday and Friday correspond to Saturday and Sunday in the United States. Thus, there was overlap on only 3 days of the workweek.

Although our research team communicated regularly by e-mail and held periodic teleconferences to discuss data and analytic issues, we found that faceto-face meetings played a critical role in resolving a number of methodologic issues, such as addressing data inconsistencies or missing data, placing analytic results in proper context, and interpreting unusual results. For example, Saudi researchers traveled to the United States to facilitate the initial analyses and to assist with data quality control procedures. In addition, the Saudis provided insight into secular trends within the Kingdom to help interpret study findings. This enabled discussion and key decision making to take place prior to construction of a final analytic data set and smoothed the process of manuscript preparation. Even though much work on the manuscript was accomplished by e-mail and telephone communications, it was still important for the whole team to travel to a central location and meet to finalize manuscripts for publication.

SANG HEALTH CARE SYSTEM

King Abdul Aziz Medical City is a tertiary care referral center in Riyadh, including a 560-bed hospital. Universal health care is provided to SANG soldiers and their family members, as well as military retirees. Each hospital visit generates an electronic record with a sequential medical record number associated with a unique SANG identification number. The SANG identification number is used for health care eligibility by SANG family members and sometimes by extended family members, but each patient has his own individual medical record. By introducing a system of taxonomy for soldiers or family members into the automated records, we were able to distinguish the categories.

Although vital statistics reporting is mandatory in Saudi Arabia, these data presented some challenges. Cause of death is usually recorded as "cardio-respiratory arrest," and autopsies are generally not conducted. Date of birth information is not based on the Gregorian calendar but instead on the lunar year. Also, by custom, if the date of birth is unknown, Saudi citizens use a fixed day and month based on royal decree.

The majority of SANG troops are Bedouin, traditionally nomadic desert dwellers, serving as full-time

domestic security forces in local geographic regions, and the vast majority are likely to remain in the SANG over their entire career. In general, SANG soldiers are characterized as stoic and not likely to seek health care for minor complaints. Differences in how SANG soldiers perceive health and illness and the impact of this perception on health care-seeking behaviors likely affected utilization patterns. In fact, speculation concerning possible adverse effects of the 1991 Gulf War was not a priority among the Saudi health care providers or the SANG military leadership, and the notion that illnesses in Gulf War veterans might well represent a condition unique to English-speaking peoples was mentioned by SANG health care providers more than once during informal discussions.

As is the case with US health care institutions, Saudi health care institutions also generally regard the sharing of individual level health data as potentially problematic. Therefore, adequately addressing patient confidentiality issues assumed the highest priority. We obtained approvals from all appropriate institutional review boards (USUHS, KAMC, CDC, and NHRC) and took measures to ensure high-level support for our project before data collection was initiated. The endorsement of the former Chief Executive Officer of Saudi Arabian National Guard Health Affairs was critical. The data team devised a process to conduct data linkages using unique study identifiers to maintain individual record integrity without retaining personal identifying information as part of the final analytic data set.

CONCLUSIONS

To our knowledge, this endeavor was the first epidemiologic investigation of postwar health outcomes among coalition forces native to the Gulf region following the 1991 Gulf War. In addition to methodologic issues, the members of the Saudi-US team of scientists had to remain aware of and sensitive to the differences between our 2 societies, which impacted all phases of our research collaboration. By paying particular attention to distinct social and religious customs, geopolitical differences, and unique aspects of the health care system, we achieved a successful international collaboration in health research.

In addition to all that we learned about the process of gathering information on health outcomes in an international setting, we gained much in the area of improved cross-cultural understanding and cooperation, which will benefit both Saudi and US scientists engaged in future health research. Despite the many challenges, the foundation for future Saudi-US scientific collaboration was established, including the potential for establishing a broad-based, multinational health surveillance system for the mutual benefit of

all participants. The Middle East in general, and the Arabian Gulf region in particular, take on added significance, given current political events and the likelihood of continued multinational operations, military and/or humanitarian, in that area of the world. Our experience contributes toward building a global network of culturally competent scientists committed to population health research. We recommend that future opportunities for international collaborative research be encouraged and supported.

The authors thank the following individuals, including their staffs, for all their support, which was critical to the success of our research collaboration: HRH General Mit'eb Bin Abdullah Bin Abdulaziz, Assistant Deputy Commander, SANG; Chief Executive Officer, Dr. Abdullah Al Rabeeah, SANG-Health Affairs; and former Chief Executive Officer, Dr. Fahad Bin Abdullah Al Abduljabbar, SANG-Health Affairs; and the SANG military leadership and the staff at KAMC for their support and Dr. Michael E. Kilpatrick from the Deployment Health Support Directorate (formerly Office of Special Assistant for Gulf War Illnesses) for his assistance with establishing this collaboration.

References

- Gackstetter GD, Hooper TI, Al Qahtani MS, Smith TC, Memish ZA, Schlangen KM, et al. Assessing the potential health impact of the 1991 Gulf War on Saudi Arabian National Guard soldiers. Int J Epidemiol 2005. In press.
- The Research Working Group of the Persian Gulf Veterans Coordinating Board. Annual report to Congress: federally sponsored research on Gulf War veterans' illnesses for 2001. June 2002. Available from: http://wwwl.va.gov/resdev/resources/pubs/docs/gulf_ war_report_2000.pdf. Accessed January 7, 2004.
- Goss Gilroy Inc. Health Study of Canadian Forces Personnel Involved in the 1991 Conflict in the Persian Gulf. Vol I. Ottawa, Ontario: Goss Gilroy Inc; 1998. Available from: http://www.forces.gc.ca/site/reports/ Health/vol I_TOC_e.htm. Accessed January 7, 2004.
- Coker WJ, Bhatt BM, Blatchley NF, Graham JT. Clinical findings for the first 1000 Gulf war veterans in the Ministry of Defence's medical assessment programme. BMJ 1999;318:290-4.
- Sim M, Abramson M, Forbes A, Glass D, Ikin J, Ittak P, et al. Australian Gulf War Veterans' Health Study. Commonwealth Department of Veterans' Affairs, Australia. Available from: http://www.dva.gov.au/ media/publicat/2003/gulfwarhs/html/prelim.htm. Accessed January 7, 2004
- Barrett DH, Gray GC, Doebbeling BN, Clauw DJ, Reeves WC. Prevalence of symptoms and symptom-based conditions among Gulf War veterans: current status of research findings. Epidemiol Rev 2002; 24:218-27.
- Doebbeling BN, Woolson RF, Torner JC, Merchant JA, Barrett DH, Polzer JP, et al. Self-reported illness following service in the Persian Gulf: multiple medical conditions. Annual Meeting of the American Public Health Association, Indianapolis, IN, November 1997 (abstract).
- Doebbeling BN, Clarke WR, Watson D, Torner JC, Woolson RF, Voelker MD, et al. Is there a Persian Gulf War syndrome? Evidence from a large population-based survey of veterans and nondeployed controls. Am J Med 2000;108:695-704.
- Fukuda K, Nisenbaum R, Stewart G, Thompson WW, Robin L, Washko RM, et al. Chronic multisymptom illness affecting Air Force veterans of the Gulf War. JAMA 1998;280:981-8.
- Gray GC, Coate BD, Anderson CM, Kang HK, Berg SW, Wignall FS, et al. The postwar hospitalization experience of US veterans of the Persian Gulf War. N Engl J Med 1996;335:1505-13.
- Gray GC, Hawksworth A, Smith TC, Kang HK, Knoke JD, Gackstetter GD. Gulf War veterans' health registries. Who is most likely to seek evaluation? Am J Epidemiol 1998;148:343-9.

- Gray GC, Kaiser KS, Hawksworth AW, Hall FW, Barrett-Connor E. Increased postwar symptoms and psychological morbidity among US Navy Gulf War veterans. Am J Trop Med Hyg 1999;60:758-66.
- Gray GC, Smith TC, Knoke JD, Heller JM. The postwar hospitalization experience of Gulf War Veterans possibly exposed to chemical munitions destruction at Khamisiyah, Iraq. Am J Epidemiol 1999;150:532-40.
- Gray GC, Smith TC, Kang HK, Knoke JD. Are Gulf War veterans suffering war-related illnesses? Federal and civilian hospitalizations examined, June 1991 to December 1994. Am | Epidemiol 2000;151:63-71.
- Gray GC, Reed RJ, Kaiser KS, Smith TC, Gastanaga VM. Self-reported symptoms and medical conditions among 11,868 Gulf War-era veterans: the Seabee Health Study. Am J Epidemiol 2002;155:1033-44.
- Gray G, Kaiser K, Hawksworth AW, Watson HL. No serologic evidence that Gulf War veterans are suffering postwar morbidity from Mycoplasma fermentans infection. Am | Trop Med Hyg 1999;60:752-7.
- Horner RD, Kamins KG, Feussner JR, Grambow SC, Hoff-Lindquist J, Harati Y, et al. Occurrence of amyotrophic lateral sclerosis among Gulf War veterans. Neurology 2003;61:743-9.
- Hyams KC, Wignall FS, Roswell R. War syndromes and their evaluation: from the US Civil War to the Persian Gulf War. Ann Intern Med 1996;125:398-405.
- Hyams KC. Gulf War syndrome: potential role of infectious diseases. Curr Opin Infect Dis 1999;12:439-43.
- Hyams KC, Riddle J, Trump DH, Wallace MR. Protecting the health of United States military forces in Afghanistan: applying lessons learned since the Gulf War. Clin Infect Dis 2002;34(Suppl 5):S208-14.
- Kipen HM, Hallman W, Kang H, Fiedler N, Natelson BH. Prevalence of chronic fatigue and chemical sensitivities in Gulf Registry veterans. Arch Environ Health 1999;54:313-8.
- Kang HK, Bullman T. Mortality among US veterans of the Persian Gulf War. N Engl | Med 1996;335:1498-504.
- Kang HK, Mahan CM, Lee KY, Magee CA, Murphy FM. Illnesses among United States veterans of the Gulf War: a population-based survey of 30,000 veterans. J Occup Environ Med 2000;42:491-501.
- Kang HK, Bullman TA, Macfarlane GJ, Gray GC. Mortality among US and UK veterans of the Persian Gulf War: a review. Occup Environ Med 2002;59:794-9.
- Kang HK, Mahan CM, Lee KY, Murphy FM, Simmens SJ, Young HA, et al. Evidence for a deployment-related Gulf War syndrome by factor analysis. Arch Environ Health 2002;57:61-8.
- Kang HK, Natelson BH, Mahan CM, Lee KY, Murphy FM. Posttraumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30,000 veterans. Am J Epidemiol 2003;157:141-8.
- Nisenbaum R, Barrett D, Reyes M, Reeves W. Deployment stressors and a chronic multisymptom illness among Gulf War veterans. J Nerv Ment Dis 2000;188:259-66.
- Riddle JR, Brown M, Smith T, Ritchie EC, Brix KA, Romano J. Chemical warfare and the Gulf War: a review of the impact on Gulf veterans' health. Mil Med 2003;168:606-13.
- Smith TC, Gray GC, Knoke JD. Is systemic lupus erythematosus, amyotrophic lateral sclerosis, or fibromyalgia associated with Persian Gulf War service? Examination of Department of Defense hospitalization data. Am J Epidemiol 2000;151:1053-9.
- Smith TC, Heller JM, Hooper TI, Gackstetter GD, Gray GC. Are Gulf War veterans experiencing illness due to exposure to smoke from Kuwaiti oil well fires? Examination of Department of Defense hospitalization data. Am J Epidemiol 2002;155:908-17.
- Smith TC, Smith B, Ryan MA, Gray GC, Hooper TI, Heller JM, et al. Ten years and 100,000 participants later: occupational and other factors influencing participation in US Gulf War health registries. J Occup Environ Med 2002;44:758-68.
- 32. Smith TC, Gray GC, Weir JC, Heller JM, Ryan MAK. Gulf War veterans and Iraqi nerve agents at Khamisiyah: postwar hospitalization data revisited. Am J Epidemiol 2003;158:456-67.

196 Vol. 33 No. 3

- Smith TC, Jimenez DL, Smith B, Gray GC, Hooper TI, Gackstetter GD, et al. The postwar hospitalization experience of Gulf War veterans participating in US health registries. J Occup Environ Med 2004;46:386-97.
- Zwerling C, Torner JC, Clarke WR, Voelker MD, Doebbeling BN, Barrett DH, et al. Self-reported postwar injuries among Gulf War veterans. Public Health Rep 2000;115:346-9.
- Unwin C, Blatchley N, Coker W, Ferry S, Hotopf M, Hull L, et al. Health of UK servicemen who served in the Persian Gulf War. Lancet 1999;353:169-82.
- Steele L. Prevalence and patterns of Gulf War illness in Kansas veterans: association of symptoms with characteristics of person, place, and time of military service. Am J Epidemiol 2000;152:992-1002.
- Cherry N, Creed F, Silman A, Dunn G, Baxter D, Smedley J, et al. Health and exposures of United Kingdom Gulf War veterans. Part I: The pattern and extent of ill health (see comment). Occup Environ Med 2001;58:291-8.
- Hallman WK, Kipen HM, Diefenbach M, Boyd K, Kang H, Leventhal H, et al. Symptom patterns among Gulf War registry veterans. Am J Public Health 2003;93:624-30.



Send us your new address at least six weeks ahead

Don't miss a single issue of the journal! To ensure prompt service when you change your address, please photocopy and complete the form below.

Please send your change of address notification at least six weeks before your move to ensure continued service. We regret we cannot guarantee replacement of issues missed due to late notification.

JOURNAL TITLE: Fill in the title of the journal here.		
OLD ADDRESS: Affix the address label from a recent issue of the journal here.	NEW ADDRESS: Clearly print your new address here. Name Address	
	City/State/ZIP	

COPY AND MAIL THIS FORM TO:

Subscription Customer Services Elsevier Inc. 6277 Sea Harbor Dr Orlando, FL 32887 OR FAX TO: 407-363-9661 OR E-MAIL: elspcs@elsevier.com **OR PHONE:** 800-654-2452 Outside the U.S., call 407-345-4000

REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

3. DATES COVERED (from - to) 1. Report Date (DD MM YY) 2. Report Type June 20, 2000 New October 1995 4. TITLE AND SUBTITLE 5a. Contract Number: Saudi Arabia-United States Collaboration in Health Research: A Formula for 5b. Grant Number: Success 5c. Program Element: 63A 5d. Project Number: **6. AUTHORS** T. I. Hooper, T. C. Smith, G. C. Gray, M. S. Al Qahtani, Z.A. Memish, D. H. 5e. Task Number: Barrett, K. M. Schlangen, D. F. Cruess, M. A. K. Ryan 5f. Work Unit Number: 60002 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P.O. Box 85122 San Diego, CA 92186-5122 9. PERFORMING ORGANIZATION REPORT **NUMBER** 8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES) Report No. 04-12 Chief, Bureau of Medicine and Surgery 10. Sponsor/Monitor's Acronyms(s) 2600 E. St., NW Washington, DC 20372-5300 11. Sponsor/Monitor's Report Number(s)

12 DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT (maximum 200 words)

Background International collaborations offer new perspectives and important opportunities for health researchers to share information, experiences, and methods that provide the basis for population-based health studies of mutual interest. Collaborations that involve completely different cultural perspectives also present distinct challenges. We share our experiences following a successful international collaboration to study health outcomes among Saudi Arabian National Guard (SANG) soldiers following the 1991 Gulf War. **Methods** Our study involved a multidisciplinary team, including two physician researchers from the Kingdom of Saudi Arabia and eight researchers from the United States with expertise in epidemiology, preventive medicine, occupational medicine, biostatistics, data programming, and analyses.

Results The findings from our epidemiological study are reported separately. In this commentary, we address some of the major distinctions between the Kingdom of Saudi Arabia and the United States, including geopolitical, social and religious, as well as healthcare system and language differences that had an impact on our research collaboration.

Conclusions Despite the challenges, our international collaboration was a very worthwhile experience. In addition to all that was learned about the process of gathering information on health outcomes in an international setting, we also gained much in the area of improved cross-cultural understanding and cooperation, which can contribute to a global network of scientists committed to building a science-based foundation for health research

15. SUBJECT TERMS

collaboration, international cooperation, Saudi Arabia, Persian Gulf syndrome, Gulf War syndrome, morbidity, healthcare, military personnel, military medicine, military deployment

16. SECURITY CLASSIFICATION OF:			17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON
		b. THIS PAGE	OF ABSTRACT UNCL	OF PAGES	Commanding Officer
UNCL	UNCL	UNCL	ONCL	5	19b. TELEPHONE NUMBER (INCLUDING AREA CODE)
					COMM/DSN: (619) 553-8429